

University of Groningen

Biochemical markers predictive for bone marrow involvement in systemic mastocytosis

Donker, Marjolein L.; van Doormaal, Jasper J.; van Doormaal, Frederiek F.; Kluin, Philip M.; van der Veer, Eveline; de Monchy, Jan G. R.; Kema, Ido P.; Kluin-Nelemans, Hanneke C.

Published in:
Haematologica-the Hematology Journal

DOI:
[10.3324/haematol.11558](https://doi.org/10.3324/haematol.11558)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2008

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Donker, M. L., van Doormaal, J. J., van Doormaal, F. F., Kluin, P. M., van der Veer, E., de Monchy, J. G. R., Kema, I. P., & Kluin-Nelemans, H. C. (2008). Biochemical markers predictive for bone marrow involvement in systemic mastocytosis. *Haematologica-the Hematology Journal*, 93(1), 120-123.
<https://doi.org/10.3324/haematol.11558>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Biochemical markers predictive for bone marrow involvement in systemic mastocytosis

Marjolein L. Donker,^{1,2} Jasper J. van Doormaal,² Frederiek F. van Doormaal,² Philip M. Kluin,³ Eveline van der Veer,⁴ Jan G.R. de Monchy,² Ido P. Kema,⁴ and Hanneke C. Kluin-Nelemans¹

Departments of Internal Medicine, ¹Hematology and ²Allergic diseases, ³Pathology and ⁴Laboratory Medicine, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

ABSTRACT

Systemic mastocytosis is characterized by bone marrow involvement, which requires a bone marrow biopsy for diagnostic work-up. We questioned whether bone marrow involvement could be predicted using biochemical markers. We selected patients with various symptoms suggestive of indolent systemic mastocytosis, of whom 63 ultimately had bone marrow involvement. Patients suspected of aggressive mastocytosis, or mastocytosis associated with other hematologic diseases were excluded. Evaluation of 115 patients and 15 patient controls demonstrated a test accuracy for serum tryptase, urinary N⁺ methylhistamine and N⁺ methylimidazole acetic acid of 96%, 88% and 95% respectively. These markers provide an excellent pre-test probability of indolent systemic mastocytosis.

Key words: mastocytosis, bone marrow involvement, biochemical markers, tryptase, N⁺ methylhistamine, N⁺ methylimidazole acetic acid.

Citation: Donker ML, van Doormaal JJ, van Doormaal FF, Kluin PM, van der Veer E, de Monchy JGR, Kema IP, Kluin-Nelemans HC. Biochemical markers predictive for bone marrow involvement in systemic mastocytosis. *Haematologica*. 2008 Jan; 93(1):120-123. DOI: 10.3324/haematol.11558

©2008 Ferrata Storti Foundation. This is an open-access paper.

Introduction

Mastocytosis denotes a heterogeneous group of disorders characterized by abnormal growth and accumulation of mast cells in one or more organ systems. In systemic mastocytosis the bone marrow is always involved.¹ A bone marrow biopsy with appropriate immunohistochemical studies is considered the gold standard for diagnosis. Ancillary tests such as mast cell immunophenotyping, cytogenetic/molecular studies, and serum tryptase levels assist in confirming the diagnosis.² While patients with aggressive systemic mastocytosis and mast cell leukemia are usually severely symptomatic, adult patients with indolent mastocytosis frequently present only with skin lesions and/or mediator-related symptoms.³ As these patients may be reluctant to undergo a bone marrow biopsy, it might be helpful if biochemical parameters could select, with a high level of certainty, patients with and without bone marrow involvement. Analysis of biochemical mediators released by mast cells may be useful for screening and diagnostic evaluation.

We investigated the predictive value and accuracy of serum

tryptase and urinary N⁺ methylhistamine (MH) and N⁺ methylimidazole acetic acid (MIMA) to predict bone marrow involvement in indolent systemic mastocytosis. We retrospectively analyzed blood, urine, and bone marrow data of a large cohort of adult patients who had been referred to our department with the probable diagnosis of indolent systemic mastocytosis.

Design and Methods

In 115 patients, a diagnosis of indolent systemic mastocytosis was considered because of: biopsy-proven cutaneous mastocytosis (all examined by an experienced dermatologist: adult-onset n=33; pediatric-onset n=3), unexplained anaphylactic reaction (n=6), hymenoptera anaphylaxis (n=22) or anaphylaxis in response to intravenous contrast material (n=1; the diagnosis of anaphylaxis was based on criteria as described in a recent paper by Sampson *et al.*⁴), mediator-related symptoms such as flushing, pruritus, abdominal pain, nausea, vomiting, diarrhea (n=16), unexplained severe osteo-

Acknowledgments: the authors thank Dr. Anneke Muller Kobold, Dr. André Mulder, Dr. Janneke Dijk-Brouwer, Mr. Henk Breukelman, and Prof. Dr. Wim Timens (Department of Pathology and Laboratory Medicine, University Medical Center, Groningen) for their expert technical assistance in immunophenotyping and measurements of KIT mutation, histamine metabolites, and for providing bone marrow biopsies of controls. We are grateful to the dermatologist Dr. Pieter van Voorst Vader, for his careful evaluation of the patients with cutaneous mastocytosis.

Funding: this study was financially supported in part by the Jan Kornelis de Cock Foundation (to FFvD), project number 564145.

Manuscript received March 27, 2007. Manuscript accepted November 6, 2007.

Correspondence: Jasper J. van Doormaal, MD, PhD, Department of Internal Medicine, Allergic Diseases, Room U3.127, University Medical Center, Groningen Hanzeplein 1, Postbox 30.001 9700 RB, Groningen, the Netherlands. E-mail: j.j.van.doormaal@int.umcg.nl

Table 1. Clinical and laboratory characteristics of subjects.

	Patients (n=115)	Controls (n=15)
Patient characteristics		
Age (years) [median, range]	49 (16-82)	54 (20-75)
Male	41 (36%)	6 (40%)
Laboratory measurements		
Tryptase (µg/L) [median, range]	22.9 (1.0-283)	4.85 (2.3-16.5)
Above 20 µg/L*	64 (56%)	
MH (µmol/mol creatinine) [median, range]	221 (44-2554)	68.5 (27-140)
Above 97 th percentile	85 (74%)	—
MIMA (mmol/mol creatinine) [median, range]	2.3 (0.5-50.6)	1.4 (0.8-2.4)
Above 97 th percentile	77 (67%)	1 (7%)
KIT mutation D816V*		
Positive	36 (31%)	—
Negative	50 (44%)	—
Not assessed	29 (25%)	15 (100%)
Immunophenotype*		
Co-expression of CD117 with CD2 and/or CD25		
Positive	39 (34%)	—
Negative	27 (23%)	—
Not assessed	49 (43%)	15 (100%)

* Minor WHO criterion¹² MH: N-methylhistamine; MIMA: N-methylimidazole acetic acid.

porosis (n=7) or combinations thereof (n=27). Patients with clinical B and C findings suggestive of aggressive mastocytosis were excluded from the study along with patients with a probable diagnosis of systemic mastocytosis with an associated hematologic non-mast cell lineage disease.² Fifteen patients who underwent a bone marrow biopsy for reasons not related to mast cell pathology served as controls. In these patients, bone marrow biopsies were performed for: analysis of competence as a bone marrow donor (n=1), cytopenia of any type (n=4), staging of non-hematologic malignancy (n=5), and exclusion of myeloproliferative disorder (n=3) or systemic amyloidosis (n=2). A bone marrow biopsy was performed and tryptase, MH and MIMA were determined for all patients and patient controls. The median interval (range) between bone marrow and blood/urine analysis was 0 (0-365) days. This time interval was acceptable if clinical symptoms remained unchanged because, in fact, biochemical parameters of patients with proven indolent mastocytosis are usually stable for several years.⁵

Tryptase levels were determined using the B12 assay.⁶ Levels of MH were determined by an isotope-dilution mass fragmentographic method.⁷ MIMA was determined as previously described⁸ with some modifications using isotope dilution mass fragmentography.

Bone marrow biopsies were taken from the iliac crest by an experienced hematologist using the Jamshidi technique. The bone marrow core required a minimal length of 20 mm to be considered representative. All

formalin-fixed paraffin-embedded (n=106) samples were stained using the Avidin-Biotin system for tryptase (clone AA1 from Dakocytomation, Glostrup, Denmark) and CD117 (Dakocytomation) expression. Tissue sections were stained by toluidin-blue for biopsies embedded in plastic (n=24).

For bone marrow mast cell immunophenotyping, 300,000 events were analyzed using four-color staining, including CD45, CD117, CD2 and CD25 as previously described.⁹ KIT mutation analysis was performed as previously described.^{10, 11}

All bone marrow biopsies were revised blinded, and scored for the presence of major and minor criteria according to the WHO classification¹² by an experienced hematopathologist (PMK). Obviously, serum tryptase as a minor criterion was omitted for this specific study.

For statistical analysis, patients were grouped into positive or negative for systemic mastocytosis by outcome of bone marrow revision, the additional minor criteria CD2 and/or CD25 expression by KIT positive mast cells, and the D816V KIT mutation.¹² Receiver operator characteristic (ROC) curves were constructed with Analyse-It (Analyse-It Software, Leeds, United Kingdom) to examine the diagnostic value of each analyte. Furthermore, we analyzed whether combinations of the three biochemical markers could increase the predictive value for bone marrow involvement compared with each marker alone.

Results and Discussion

Patient characteristics of both groups were comparable (see Table 1). Both groups consisted of comparable subjects. Results from bone marrow histology and additional WHO minor criteria are shown in Figure 1. Fifty-nine patients showed one major and one or more minor criteria for the diagnosis of systemic mastocytosis. In 51 patients, a diagnosis of systemic mastocytosis could not be made. In this retrospective study, additional data on flow cytometry or KIT mutation analysis were sometimes lacking (Table 1). However, we decided not to repeat these tests, either because there were already sufficient major and minor criteria for the diagnosis of systemic mastocytosis or because all other criteria were negative, indicating that no diagnosis of systemic mastocytosis could be made. Given that our study aim was to evaluate whether serum tryptase and the urinary metabolites MH and MIMA could predict systemic mastocytosis, all WHO major and minor criteria were used except serum tryptase. Consequently, 2 patients, whose diagnosis of systemic mastocytosis was based on this criterion (tryptase 23.9 and 28.9 µg/L, respectively), were excluded from further study (Figure 1).

ROC analyses were performed to study the sensitiv-




Figure 1. Distribution of patients according to bone marrow involvement. Distribution of subjects in indolent systemic mastocytosis/no indolent systemic mastocytosis groups used for statistical analysis, according to bone marrow pathology and evaluation of WHO minor criteria related to *KIT* mutation and CD2/CD25 immunophenotyping.

ity and specificity of serum tryptase and urinary MH and MIMA. All biochemical markers showed an area under the curve (AUC) of >80% indicating excellent test accuracy, with serum tryptase (AUC: 0.960; 95% CI: [0.930 - 0.991]) and urinary MIMA (AUC: 0.948; 95% CI: [0.911 - 0.985]) performing significantly better (tryptase vs. MH $p=0.0096$, MIMA vs. MH $p=0.0057$) than urinary MH (AUC: 0.881; 95% CI: [0.823-0.940]). There was no significant difference between diagnostic accuracy of tryptase and MIMA ($p=0.5613$). Analysis of combinations of the three biochemical markers compared with each marker alone showed no further increase in diagnostic accuracy (*data not shown*).

Although it is not necessary to choose any particular threshold for assessment of test accuracy of the various biochemical markers, we selected various thresholds to be used in patient care (Table 2). This means that a patient with only cutaneous involvement and a serum tryptase level below <20 $\mu\text{g/L}$ (threshold used by the WHO as minor criterion¹²) would have a 16% risk of bone marrow involvement, and thus a diagnosis of indolent systemic mastocytosis. This risk would further decrease to 0% if such a patient had a tryptase level <4.9 $\mu\text{g/L}$. In the absence of symptoms suggestive

of mastocytosis, and given present lack of a curative therapy for systemic mastocytosis, this could be considered acceptable. On the other hand, this could help convince patients and doctors to perform a bone marrow biopsy for those with cutaneous mastocytosis or clinical symptoms suggestive of mastocytosis who have elevated levels of serum tryptase and/or urinary histamine metabolites. The probability that a patient with a serum tryptase level of $\geq 21.9 \mu\text{g/L}$ had bone marrow involvement was 98%, and with a level of $\geq 30.8 \mu\text{g/L}$ this rose even to 100%. Notably, serum tryptase is not pathognomonic for mastocytosis. Patients with high tryptase levels may harbor another disease, such as a myeloid haematologic disorder.¹³

Remarkably, clinical symptoms were helpful in predicting bone marrow involvement. Obviously, this should be related to the referral pattern of this group of patients to our university hospital. The majority of the patients with anaphylactic reactions (14/24) and with cutaneous mastocytosis (21/31 evaluable patients) had systemic mastocytosis. In particular, patients with cutaneous mastocytosis with additional clinical symptoms were almost all classified as systemic mastocytosis (23/25). This agrees with the general assumption that

Table 2. Sensitivity/specificity pairs and predictive values. Sensitivity/specificity pairs and positive predictive values (PPV) and negative predictive values (NPV) of tryptase, MH and MIMA at various cut-off points.

Marker	Cut-off point	Sensitivity	Specificity	PPV	NPV
Tryptase	≥4.94 µg/L	100%	41.2%	67.0%	100%
	≥21.9 µg/L	83.6%	98.0%	98.4%	83.3%
	≥30.8 µg/L	67.2%	100%	100%	71.8%
MH	≥103 µmol/mol creatinine	100%	39.2%	66.3%	100%
	≥171 µmol/mol creatinine	86.9%	70.6%	77.9%	81.8%
	≥284 µmol/mol creatinine	55.7%	100%	100%	65.4%
MIMA	≥1.6 mmol/mol creatinine	100%	56.9%	73.5%	100%
	≥2.0 mmol/mol creatinine	93.4%	82.4%	86.3%	91.3%
	≥3.8 mmol/mol creatinine	49.2%	100%	100%	62.2%

adult-onset cutaneous mastocytosis commonly represents systemic mastocytosis. By contrast, 4 out of the 15 patients with mediator-related symptoms only were classified as such. The quality of the bone marrow biopsy and additional immunohistochemistry is very important, and we used strict criteria for final interpretation. Mast cell infiltrates can be sparse, therefore examination of sufficient marrow fields for a final negative diagnosis is required making additional immunohistochemistry essential. In our retrospective study, we had to discard 12% of the biopsy specimens, the majority due to small size. We used control patients instead of healthy volunteers. Besides ethical issues, the main reason for this selection was based upon the fact that the clinical question was not to identify subjects with or without bone marrow involvement in asymptomatic volunteers, but patients who are considered to have

systemic mastocytosis. In conclusion, we have demonstrated in a large representative cohort of patients that blood and urinary analysis for tryptase and histamine metabolites can help to predict those patients who have bone marrow involvement representing indolent systemic mastocytosis.

Authorship and Disclosures

MLD, JJvD, FFvD, PMK, EvdV, JGRdM, IPK, HCK-N: contributed substantially to conception and design, to acquisition of data, to analysis and interpretation of data, drafted and revised the manuscript critically for important intellectual content and gave final approval of the version to be published. The authors reported no potential conflicts of interest.

References

- Pardanani A, Akin C, Valent P. Pathogenesis, clinical features, and treatment advances in mastocytosis. *Best Pract Res Clin Haematol* 2006; 19:595-615.
- Patnaik MM, Rindos M, Kouides PA, Tefferi A, Pardanani A. Systemic mastocytosis: a concise clinical and laboratory review. *Arch Pathol Lab Med* 2007;131:784-91.
- Akin C, Metcalfe DD. Systemic mastocytosis. *Annu Rev Med* 2004; 55:419-32.
- Sampson HA, Munoz-Furlong A, Campbell RL, Adkinson NF, Jr., Bock SA, Branum A, et al. Second symposium on the definition and management of anaphylaxis: summary report--Second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network symposium. *J Allergy Clin Immunol* 2006;117:391-7.
- Kors JW, Van Doormaal JJ, Breukelman H, Van Voorst Vader PC, De Monchy JG. Long-term follow-up of indolent mastocytosis in adults. *J Intern Med* 1996;239:157-64.
- Granerus G, Lonnqvist B, Nystrand J, Roupe G. Serum tryptase measured with B12 and G5 antibody-based immunoassays in mastocytosis patients and its relation to histamine turnover. *Br J Dermatol* 1998; 139:858-61.
- Keyzer JJ, Wolthers BG, Muskiet FA, Kauffman HF, Groen A. Determination of N tau-methylhistamine in plasma and urine by isotope dilution mass fragmentography. *Clin Chim Acta* 1981;113:165-73.
- Keyzer JJ, Wolthers BG, Breukelman H, Kauffman HF, de Monchy JG. Determination of N tau-methylimidazoleacetic acid (a histamine metabolite) in urine by gas chromatography using nitrogen-phosphorus detection. *Clin Chim Acta* 1982;121:379-87.
- Pardanani A, Kimlinger T, Reeder T, Li CY, Tefferi A. Bone marrow mast cell immunophenotyping in adults with mast cell disease: a prospective study of 33 patients. *Leuk Res* 2004;28:777-83.
- Nagata H, Worobec AS, Oh CK, Chowdhury BA, Tannenbaum S, Suzuki Y, et al. Identification of a point mutation in the catalytic domain of the protooncogene c-kit in peripheral blood mononuclear cells of patients who have mastocytosis with an associated hematologic disorder. *Proc Natl Acad Sci USA* 1995;92:10560-4.
- Worobec AS, Semere T, Nagata H, Metcalfe DD. Clinical correlates of the presence of the Asp816Val c-kit mutation in the peripheral blood mononuclear cells of patients with mastocytosis. *Cancer* 1998;83:2120-9.
- Valent P, Horny HP, Li CY, Longley JB, Metcalfe DD, Parwaresch RM, et al. Mastocytosis (mast cell disease). World Health Organization (WHO) classification of tumours. Pathology and genetics. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, editors. *Tumours of haematopoietic and lymphoid tissues*. Vol 1. World Health Organization. Lyon, France: IARC Press, 2001;291-302.
- Schwartz LB. Clinical utility of tryptase levels in systemic mastocytosis and associated hematologic disorders. *Leuk Res* 2001;25:553-62.